



<h1 style="margin: 0;">AE3600</h1> <p style="margin: 10px 0 0 0;">3.6 RM</p>	Trait Options	Geographic Adaptation				PRO-FIT Placement				
	Xtend ■	Western West of Miss. River	HR	Moves South well for early season variety	HR	Outstanding disease tolerance in tough locations				
	Enlist ■	Central (I-35 to I-65)	HR		Combination of offense and defense					
	LL/GT27 ■	Eastern East of Miss. River	HR	Moves North well for full season variety	HR	Dependable standability				
Conventional ■					Performs from Missouri to Ohio					
General Characteristics			Agronomic Features				Disease Tolerance			
Plant Height	Medium	Yield for Maturity	Very Good		Brown Stem Rot	Susceptible				
Canopy/Plant type	Thin	Emergence	Very Good		Frogeye Leaf Spot	NA				
Growth Habit	Upright	Standability	Excellent		Iron Chlorosis	Fair				
Flower Color	White	Shatter Resistance	Very Good		Phytophthora Field Tol.	Very Good				
Hilum Color	Buff	Stress Tolerance	Very Good		SDS Tolerance	Good				
Pod Color	Brown	SCN Resistance Source	PI88788		White Mold	NA				
Pubescence	Gray	Phytophthora Res. Source	None		Stem Canker	NA				
Protein	NA	No-Till Adaptability	Very Good		Root Knot	NA				
Oil	NA					Cover Crop Companion	Very Good			
Soil Placement		Recommended Seeding Rates				Herbicide Tolerance				
Light Soils: Low O.M., Low CEC, Low water holding capacity.	HR	Row Width	120/130	135/145	150/160	Conventional	■	Dicamba	■	
Medium Soils: O.M. 1.5-3.5%, CEC 11-20, Good productivity, well drained silt loams.	HR	30" Rows	HR	HR	R	Glyphosate	■	2,4-D	■	
Heavy Soils: High O.M. >3.5%, CEC >20, Well drained highly productive with deep topsoil	HR	Twin Rows	HR	HR	R	Glufosinate	■	Callisto	■	
Poorly Drained Soils: Soils that tend to remain saturated for extended periods of time.	HR	20" Rows	NR	R	HR	Sulfonlurea	■	IsoxoflutoI	■	
		<20" rows	NR	R	HR					
Population ranges are listed in 1000 seeds/acre										
NR=Not Recommended, R=Recommended, HR=Highly Recommended										
Notes:										